

**SEVERE COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hci liquid
P & L Development, LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- runny nose and sneezing
- cough to help you sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or last more than 3 days
- redness or swelling is present
- cough comes back, or occurs with rash or headache that lasts.

These could be a signs of a serious conditions.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **take only as directed - see Overdose warning**
- mL = milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- do not exceed 4 doses per 24 hours
- adults and children 12 years and over: 30 mL every 4 hours
- children under 12 years of age: do not use
- **When using Day Time or Night Time products, carefully read each label to ensure correct dosing.**

Other information

- **each 30 mL contains;** sodium 79 mg
- store between 15-30°C (59-86°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, FD&C blue1, FD&C red 40, Flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to active ingredients in Vicks® NyQuil® Severe Cold & Flu*

maximum strength

severe night time cold & flu

cold & flu

Acetaminophen

dextromethorphan HBr

doxylamine succinate

phenylephrine HCl

relieves:

- aches, fever & sore throat
- cough
- runny nose & sneezing
- nasal & sinus Congestion

alcohol free

berry flavor

FL OZ (mL)

*This product is not manufactured or distributed by Procter & Gamble, distributor or Vicks® NyQuil® Severe Cold & Flu.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL IS BROKEN OR MISSING.

Manufactured by:

PL Developments

11865 S. Alameda St

Lynwood, CA 90262

Product Label

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TAMPER EVIDENT: DO NOT USE IF PRINTED
SAFETY SEAL IS BROKEN OR MISSING.

PLD-A341A LB003625

*This product is not manufactured or distributed
by Procter & Gamble, distributor of Vicks®
NyQuil® Severe Cold & Flu.



Manufactured by:
PL Development
11865 S. Alameda St
Lynwood, CA 90262



3 49580 41608 6



Compare to active ingredients in
**Vicks® NyQuil® Severe
Cold & Flu***
NDC 49580-0416-8

maximum strength
**severe
night time
cold & flu**
Acetaminophen
dextromethorphan HBr
doxylamine succinate
phenylephrine HCl

relieves:

- aches, fever & sore throat
- cough
- runny nose & sneezing
- nasal & sinus congestion

alcohol free

8 fl oz (237 mL)



berry flavor

Drug Facts

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses • temporarily relieves common cold/flu symptoms:
• nasal congestion • sinus congestion and pressure
• cough due to minor throat and bronchial irritation • minor aches and pains • headache • fever • sore throat
• reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • runny nose and sneezing • cough to help you sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 4,000 mg in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued under label) **PEEL HERE** ➔

Drug Facts (continued)

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema • a sodium- restricted diet

Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin

When using this product • do not take more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if • nervousness, dizziness, or sleeplessness occur • pain, nasal congestion, or cough gets worse or lasts more than 7 days • new symptoms occur • fever gets worse or lasts more than 3 days • redness or swelling is present • cough comes back, or occurs with rash or headache that lasts.

Drug Facts (continued)

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions • take only as directed - see Overdose warning • mL = milliliter • use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device. • do not exceed 4 doses per 24 hours
• adults and children 12 years and over: 30 mL every 4 hours
• children under 12 years of age: do not use • when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information • each 30 mL contains: sodium 79 mg • store between 15-30°C (59-86°F) • do not refrigerate

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Readyincase NightTime Severe Cold & Flu Berry Liquid

SEVERE COLD AND FLU NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hci liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49580-0416
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
		650 mg

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0416-8	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/31/2015	

Labeler - P & L Development, LLC (101896231)